



### Census Results:

Conducted two population-wide censuses of nests: one in the incubation period (mid-July) and one after the hatching period (mid-September).

Census Month	Number Adults	Number Eggs	Number Chicks	Active Nests	Abandoned Eggs	Lost Eggs	Dead Chicks	Lost Nests
July	205	21	0	226	0	37	0	37
Sept.	10	0	81	91	0	17	0	17

<p><b>July Incubation Rate:</b> 87.9% (175 of 199 adults incubating, 6 unknown)</p>	<p><b>Sept. Incubation Rate:</b> 0% (0 of 9 adult incubating, 1 unknown)</p>
<p><b>Adults</b> refers to the number of nests where one or two adult birds present, but an egg not readily visible.  <b>Lost Eggs</b> refers to predated or broken eggs in a nest. <b>Abandoned Eggs</b> were found outside of a nest.  <b>Active Nests</b> includes nests occupied by one or two adults (whether incubating or not), an egg or a chick.  <b>Lost Nests</b> includes abandoned, predated and broken eggs that will not hatch, and dead chicks.</p>	

### Census Results:

Monitored 60 nests and recorded the hatching dates and the growth of 35 chicks from mid July to mid Nov.

Hatching dates:

Aug. 3 - 15.

Mean = Aug 7 (n = 30)

